



A



B



CHO cells were transfected in a Costar® 24-well tissue culture treated plate using the magnefect-nano™ transfection system (24-well plate magnet) with 0.6 µl nTMAG and 0.6 µg of a plasmid encoding EGFP per well. 48 hours post-transfection, the cells were analysed by light (A) and by fluorescence microscopy (B).

Grey (marked with a ●) denotes values for the 96-well plate magnet format

Blue (marked with a ■) denotes values for the 24-well plate magnet format

Cell seeding parameters

Plate type: ● Greiner Bio-One 96-well or ■ Costar® 24-well plate (tissue culture treated)

Cell seeding density: ● 1.5×10^4 or ■ 9.0×10^4 cells/well (4.7×10^4 cells/cm²)

Cell seeding volume: ● 100 µl or ■ 600 µl

Cell seeding time: 24 hours prior to transfection

Serum starve: **Yes; remove cell seeding medium 16 hours prior to transfection, replace with ● 100 µl or ■ 600 µl Ham's F12 Medium (i.e. serum- and supplement-free)**

Transfection complex (see magnefect-nano™ protocol for more details)

Transfection reagent: nTMAG

Transfection reagent (volume/well): ● 0.1 µl or ■ 0.6 µl

DNA (mass/well): ● 0.1 µg or ■ 0.6 µg

Transfection medium: Ham's F12 Medium (i.e. serum- and supplement-free)

Transfection volume: ● 100 µl or ■ 600 µl

Medium change after transfection: **Yes; 6 hours post-transfection replace medium with ● 100 µl or ■ 600 µl Ham's F12 Medium; 10% FCS; L-glutamine; antibiotics**

magnefect-nano™ parameters

Frequency: 2 Hz

Displacement: 0.2 mm

Time: 1 hour (7200 cycles)

Cell information

Cell type: CHO (cell line; adherent)

Species: Chinese Hamster

Tissue origin: Ovary

Morphology: Epithelial

Medium: Ham's F12 Medium; 10% FCS; L-glutamine; antibiotics

Culture condition: Temperature, 37°C; Atmosphere, 95% air, 5% CO₂