



## MG63

A



B



MG63 cells were transfected in a Greiner Bio-One 96-well tissue culture treated plate using the magnefect-nano™ transfection system (96-well plate magnet) with 0.1 µl nTMAG and 0.1 µg of a plasmid encoding EGFP per well. 48 hours post-transfection, the cells were analysed by light (A) and by fluorescence microscopy (B).

Grey (marked with a ●) denotes values for the 96-well plate magnet format

Blue (marked with a ■) denotes values for the 24-well plate magnet format

### Cell seeding parameters

Plate type: ● Greiner Bio-One 96-well or ■ Costar® 24-well plate (tissue culture treated)

Cell seeding density: ●  $1.5 \times 10^4$  or ■  $9.0 \times 10^4$  cells/well ( $4.7 \times 10^4$  cells/cm<sup>2</sup>)

Cell seeding volume: ● 100 µl or ■ 600 µl

Cell seeding time: 24 hours prior to transfection

Serum starve: No

### Transfection complex (see magnefect-nano™ protocol for more details)

Transfection reagent: nTMAG

Transfection reagent (volume/well): ● 0.1 µl or ■ 0.6 µl

DNA (mass/well): ● 0.1 µg or ■ 0.6 µg

Transfection medium: DMEM Medium (i.e. serum- and supplement-containing)

Transfection volume: ● 100 µl or ■ 600 µl

Medium change after transfection: No

### magnefect-nano™ parameters

Frequency: 2 Hz

Displacement: 0.2 mm

Time: 1 hour (7200 cycles)

### Cell information

Cell type: MG63 (cell line; adherent)

Species: Human

Tissue origin: Bone

Morphology: Fibroblast

Medium: DMEM Medium; 10% FCS; L-glutamine; antibiotics

Culture condition: Temperature, 37°C; Atmosphere, 95% air, 5% CO<sub>2</sub>